



Efficacy of Trans-Mastoidal Vestibular Galvanic Stimulation in Improvement of Gait Performance and Upright Postural Stability in Hemiplegic C.P. Children

Ahmed M. Azzam^{1*}

¹*Department of Physiotherapy for Developmental Disturbance and Pediatric Surgery,
Faculty of Physical Therapy, Cairo University, Giza, Egypt.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/CJAST/2019/v34i630149

Editor(s):

(1) Dr. Teresa De Pilli, Assistant Professor, Department of Science of Agriculture of Food of Environment (SAFE), University of Foggia, Via Napoli, Italy.

Reviewers:

(1) David Vance, University of Alabama at Birmingham, USA.

(2) Hank White, Shriners Hospitals for Children, USA.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48264>

Received 27 November 2018

Accepted 08 April 2019

Published 19 April 2019

Original Research Article

ABSTRACT

Objectives: This work was carried out to investigate the efficacy of galvanic vestibular stimulation in improving of gait performance and upright postural stability in hemiplegic cerebral palsy children.

Method: Thirty children were enrolled in this study and randomly assigned into two groups; group A (galvanic vestibular stimulation plus vestibular training program), and group B (vestibular training program only). Stride length and time, walking speed tests and modified Ashworth, pediatric balance scales were used to detect and follow the walking performance and upright postural stability. This measurement was taken before initial treatment and after 12 weeks of treatment. The children parents in both groups A and B were instructed to complete 3 hours of the home routine program.

Results: Data analysis was available on the 30 hemiplegic cerebral palsied children participated in the study. The difference between pre and post-treatment results was significant representative in stride length, time and walking velocity, spasticity changes and pediatric balance scores in study groups while insignificant improvement in control groups.

Conclusion: The combined vestibular training program and trans-mastoidal vestibular galvanic

*Corresponding author: E-mail: Prof.ahmedazzam@yahoo.com;

stimulation are suggested in improving walking performance and upright postural stability in a static and dynamic situation. So this selective physiotherapy approach may be used as a strong choice for improving walking and balance abilities in hemiplegic C.P children.

Keywords: Galvanic vestibular stimulation; cerebral palsy; walking performance.

1. INTRODUCTION

Central nervous system lesion which occurs in cerebral palsy patients lead to impaired postural reaction mechanism, loss of reciprocal inhibition mechanism, muscle tone disturbance, impaired co-ordination mechanism and failure of posture stabilization. Poor balance mechanism is the direct source of the defective motor and functional skills acquisition and motor development delay [1,2,3].

Learning process through sensory feedback is the plasticity way for continuous modifications of neural connections. Environmental manipulation and external stimulation reshape the neural network patterns through synaptic competition mechanism with the greatest activated synaptic connections become the winner and gain the skill and the lowest activated become the loser. Throughout of life there is an alteration of the synapses functional characteristics according to the new experience and skills plus environmental elements and variant situations or occurring CNS lesion [4]. Brain plasticity can occur at every lifetime because the CNS is continually remodeling via synaptic reorganization so vestibular restoration therapy rely on the active participation which produces anatomical and permanent changes in the synaptic network of the neural circuits which produce behavioral changes on the postural stability [5,6,7,8,9,10].

Utilization of vestibular galvanic electric stimulation on the mastoid process via placing of anode electrode on the mastoid process in the similar side of body sway and the cathode on the other side. This will activate Na⁺ channels opening lead to depolarization of vestibular afferent nerve which produce modulation of the afferent vestibular signals by raising firing rate of the vestibular afferent on the cathode side and reducing the excitation degree on the anodal side which leading to deviation of the posture toward anodal side [11,12,13]. GVS has a positive effect on the balance control via improve balance response time, decreasing the lateral and A-P sway in static balance together with improving dynamic balance performance and correcting the abnormal perception sway and motor response [14]. The anticipatory postural control was improved after application GVS via modulating

the vestibulospinal tract signals and activation of vestibular-related sensory areas [15].

During walking the antigravity muscle tonus (γ and α motor neurons) were modulated by the vestibulo-spinal and reticulo-spinal tracts so by the application of GVS will increase the vestibulo-spinal tract activity which transmitted to the antigravity muscles increasing their excitability leading to increasing of stride length and walking speed [16-27].

2. MATERIALS AND METHODS

2.1 Subjects

30 children from the two sexes with hemiplegic C.P. children were joined in this study, aged 5 to 10 years at a time of enrollment due to the children in this age could participate in pediatric balance scale graduations. Children could walk with assistance, On the other hand, hemiplegic C.P. Children that run up against the involvement rules were derived out if they had: preceding BoNT-A dose in the L.L or U.L in the last 12 months or had surgical tendon or muscle lengthening operation.

Children were selected randomly to the study group (A) took vestibular galvanic electric stimulation plus vestibular training approach while the control group (B) took vestibular training approach only. The individual-based vestibular galvanic electric stimulation treatment sessions of 30 minutes were conducted day after day for 3 months in a physiotherapy treatment room after the vestibular training period for the group (A). Also, children in the two groups were subjected to home regular program 3 hours daily for the 3 months treatment period. The control group (B) received a vestibular training program only.

2.2 Outcome Measurements

2.2.1 Gait speed test

It was used to evaluate the walking speed by determining a distance of 10 meters which evaluates functional vestibular abilities. Measuring tap and stopwatch tools were needed to measure the walking speed. Starting and finishing areas of one meter were used to build

up the maximum speed and decelerate of the speed. The average of 3 trials results was recorded pre and post treatment [28].

2.2.2 Modified Ashworth scale

It was used to evaluate and follow up the degree of muscle tone disturbance pre and post-treatment.

2.2.3 Pediatric balance scale

It was used to evaluate functional balance abilities. It consists of 14 items each one is scored on a 5-point scale with zero scores indicate to the child cannot achieve the task without assistance and 4 scores indicate to independence in performing the ability.

The highest score is 56 points (the summation of the scores in all items).

The points from 41-55 indicate to the independent child

The points from 21-40 indicate to the child need assistance

The points 20 or below indicate to the child need wheelchair [29,30].

2.2.4 Stride length and stride time measurements

Stride length is the extension from the toe of the foot (starting position) to the toe of same foot (ending position), or from the heel to heel of the same foot was measured by tape measurement. The stride time was detected by stopwatch. 10-meter distance (about 8 steps) was calibrated to evaluate the stride length before and after treatment. Stride time could be calculated by counting time required to perform walking in the calibrated distance.

2.3 Intervention

For all children, the treatment was handling three times weekly, for 3 months. Each session persists for 90 minutes (30 minutes vestibular galvanic electric stimulation for study group plus 60 minutes for a vestibular training program for each group) in a physical therapy room, in addition to 3 hours of the home regular program, day after day around the treatment duration

Both groups (A and B) received a vestibular training program, like the following:

1. With the eyes closed: concentration on vestibular system training through an unstable surface to isolate proprioceptors and perform postural reaction training (righting +equilibrium+ protective reaction

training) through medical balls and balance board of different sizes.

2. With the eyes opened then closed slow then rapid training. The child performed dynamic vestibular training forward to the mirror by walking sideways with feet followed each other then walked on one line then walked by passing foot to one another.
3. Static training to vestibular system by performing proximal stabilization in a quadruped, kneeling, half kneeling, standing.
4. Proprioceptive training with vestibular training by standing on one foot with eyes opened then closed.
5. Equal weight shift transfer by starting the big base of support then gradual decreased of BOS (Base of support) without disturbance then with disturbance-with eye opened then eye closed-with hand supported then without.
6. Step forward including or excluding disturbance, with eyes opened then closed on different directions and surfaces.
7. Weight-bearing with upper extremity functional training as hand skills training (from sitting-standing-kneeling) as grasping, voluntary release, reaching, hand manipulative skills, bilateral hand use, and eye-hand coordination training.
8. Changing positions from non-weight bearing to weight bearing and opposite and from static to dynamic and opposite.
9. Upside training anteroposterior movement and lateral movement then rotatory movement.
10. Swing therapy anteroposterior movement and lateral movement then rotatory movement.
11. Biodex stability system training.
12. Perturbation with different positions (quadruped-kneeling-half-kneeling-standing, holding a big medical ball with maintaining posture stability, standing against the corner, standing with holding stand bar and manual support standing).
13. Jumping training with trampoline [28].

The experimental group (group A) received vestibular galvanic electric stimulation following:

Vestibular galvanic electric stimulation is a non-invasive technique that sending a direct electric message to the vestibular receptors (3semicircular ducts and the two otolith organs

utricle and saccule) aiming for enhancement of gait performance. By locating a target to the child and starting point on a sheet and placed the anodal electrode on the mastoid process on the side of paralysis and the cathode on the other mastoid process. The elastic headbands stabilized the two stimulating electrodes. The physiotherapist asked the child to go with holding the child from his upper parts of the shoulder in the similar time of turning on the direct current with the intensity of 0.5 mA intensity with long latency more than 200 ms for 30 minutes. There is deviation of the posture on the anodal side till the child reaches to the target or the child could use a pediatric treadmill withholding of hand support and asking the child to slowly walk with the similar time of turning the apparatus on.

3. RESULTS

3.1 Patients Characteristics

Table 1 displayed the demographic and analytic traits of all patients. There were 13 boys (43.33%) and 17 girls (56.67%) and in term of right-hand dominance reported in 17 patients (56.67%), and also 13 patients (43.33%) were left-hand dominance. There was no representative change within both groups regarding age ($p=0.8816$), to sex ($p=0.7240$) and in term of hand dominance ($p=0.4814$).

3.2 Changes in Stride Length

Mean test scores and SD for both groups are demonstrated in Table 2. The mean record of stride length level in the two groups at (pre- and post-treatment levels) was worthless ($p>0.05$). The average improvement of stride length level had a tendency to be extremely representatives improvement in the experimental group (3.433 ± 0.961 versus 2.967 ± 0.990 , $p=0.0005$) while worthless representatives in the control group (3.100 ± 0.828 versus 2.967 ± 0.915 , $p=0.1038$). The percentage of improvement of stride length level was (15.706%) in the study group compared to the (4.483%) in the control group.

3.3 Changes in Walking Velocity

Mean test scores and SD for the both groups are demonstrated in Table 3. The mean record of walking velocity level in the two groups at (pre-treatment) was worthless ($p>0.05$) while the two groups had a representatives improvement in walking velocity at post-treatment level ($p<0.05$). The average improvement of walking velocity level had a tendency to be extremely representatives improvement in the study group

(6.93 ± 0.70 versus 5.8 ± 0.86 , $p=0.0001$) while worthless representatives regarding control group (5.80 ± 1.21 versus 5.67 ± 1.29 , $p=0.1643$). The percentage of improvement of walking velocity level was (19.48%) in the study group compared to the (2.29%) regarding the control group.

3.4 Changes in Pediatric Scale

Mean test scores and SD for both groups are presented in Table 4. The mean record of pediatric scale score in the two groups at pre-treatment was worthless ($p>0.05$) while the two groups had a significant improvement in pediatric scale score post-treatment ($p<0.05$). The average improvement of pediatric scale score had a tendency to be extremely representatives improvement in the study group (41.80 ± 1.32 versus 36.53 ± 3.07 , $p=0.0001$) than regarding control group (36.07 ± 2.91 versus 36.53 ± 2.72 , $p=0.0558$). The percentage of improvement of pediatric scale score was (14.43%) in the study group compared to the (1.2%) regarding control group.

3.5 Changes in Spasticity Degree

Mean test scores and SD for both groups are demonstrated in the Table 5. The mean record of spasticity degree level in the two groups at (pre-treatment) had significant improvement ($p<0.05$) while both groups had an worthless improvement in spasticity degree at post-treatment level ($p>0.05$). The average improvement of spasticity degree level tended to be extremely representatives improvement in the study group. (1.47 ± 0.52 versus 2.40 ± 0.63 , $p=0.0001$) while worthless regarding control group (1.53 ± 0.52 versus 1.60 ± 0.51 , $p=0.3343$). The percentage of improvement of spasticity degree level was (38.75%) in the study group compared to the (4.375%) regarding control group.

3.6 Changes in Stride Time

Mean test scores and SD for both groups are demonstrated in Table 6. The mean record of stride time level in the two groups at (pre-and post-treatment level) was worthless ($p>0.05$). The average improvement of stride time level had a tendency to be extremely representatives improvement in the study group (11.07 ± 1.91 versus 12.93 ± 2.25 , $p=0.0002$) than regarding control group (12.27 ± 1.58 versus 12.47 ± 1.55 , $p=0.0824$). The percentage of improvement of stride time level was (14.385%) in the study group compared to the (1.6%) regarding control group.

Table 1. Patients characteristics

Variables	Study group N=15	Control group N=15	P-value
Age	7.80±1.26	7.73±1.16	0.8816
Sex N%			
Boys	7 (46.67%)	6 (40%)	0.7240
Girls	8 (53.33%)	9 (60%)	
Hand dominance N%			
Right	9(60%)	8(53.33%)	0.4814
Left	6(40%)	7(46.67%)	

Table 2. The average test of stride length level in both groups

Stride length level	Study group Mean±SD	Control group Mean±SD	P-value (within groups)
Pre-treatment	2.967±0.990	2.967±0.915	1.000
Post-treatment	3.433±0.961	3.100±0.828	0.3176
Improvement%	15.706%	4.483%	0.0389
P-value (within groups)	0.0005	0.1038	

Table 3. The average test of walking velocity level in both groups

Walking velocity level	Study group Mean±SD	Control group Mean±SD	P-value (within groups)
Pre-treatment	5.8±0.86	5.67±1.29	0.7419
Post-treatment	6.93±0.70	5.80±1.21	0.0039
Improvement%	19.48%	2.29%	0.0004
P-value (within groups)	0.0001	0.1643	

Table 4. The average test of pediatric scale score in both groups

Pediatric scale score	Study group Mean±SD	Control group Mean±SD	P-value (within groups)
Pre-treatment	36.53±3.07	36.53±2.72	0.3531
Post-treatment	41.80±1.32	36.07±2.91	0.0001
Improvement%	14.43%	1.2%	0.3275
P-value (within groups)	0.0001	0.0558	

Table 5. The average test of spasticity degree level in both groups

Spasticity degree level	Study group Mean ±SD	Control group Mean ±SD	P-value (within groups)
Pre-treatment	2.40±0.63	1.60±0.51	0.0007
Post-treatment	1.47±0.52	1.53±0.52	0.7263
Improvement%	38.75%	4.375%	0.0009
P-value (within groups)	0.0001	0.3343	

Table 6. The average test of stride time level in both groups

Stride time level	Study group Mean ±SD	Control group Mean ±SD	P-value (within groups)
Pre-treatment	12.93±2.25	12.47±1.55	0.5140
Post-treatment	11.07±1.91	12.27±1.58	0.0710
Improvement%	14.385%	1.6%	0.0001
P-value (within groups)	0.0002	0.0824	

4. DISCUSSION

The results of the present study suggest that trans-mastoidal vestibular galvanic stimulation might be useful to enhance independency in gait performance and upright postural stability in hemiplegic C.P. children. The effects of direct electrical vestibular stimulation plus manual

vestibular training are more benefits for global function Improvement than manual vestibular training only. The study group pre and post treatment showed increase at the stride length level (3.433±0.961 versus 2.967±0.990, p=0.0005) while insignificant in the control group (3.100±0.828 versus 2.967±0.915, p=0.1038). Also the study group pre and post treatment

showed increase at the walking velocity (6.93±0.70 versus 5.8±0.86, p=0.0001) while worthless representatives regarding control group (5.80±1.21 versus 5.67±1.29, p=0.1643). In addition to the study group pre and post treatment showed increase of the of pediatric scale score (41.80±1.32 versus 36.53±3.07, p=0.0001) while insignificance in control group (36.07±2.91 versus 36.53±2.72, p=0.0558), the study group pre and post treatment showed decrease of the spasticity degree (1.47±0.52 versus 2.40±0.63, p=0.0001) while worthless regarding control group (1.53±0.52 versus 1.60±0.51, p= 0.3343) and the study group pre and post treatment showed increase of stride time level (11.07±1.91 versus 12.93±2.25, p=0.0002) while insignificance regarding control group (12.27±1.58 versus 12.47±1.55, p= 0.0824).

Utricule and saccule stimulation occurs by placing galvanic stimulation on mastoid processes in which cathode on one side and anode on another side which lead to sway on the anodal side. The perception tilt sway occur toward the paralyzed side so the anode should be located on the mastoid process of the paralyzed side because it decreases the firing rates in vestibular afferent of anodal current this will cause shifting

in subject perception sway(tilt) in a direction opposite to that turned in walking [13,31].

When a patient walks in a well-learned place he will have a pre-programmed strategy and neural motor circuits so when the patient makes a fast walking there is decrease in sway as a result of vestibular signals is less vital in fast walking as it depends on pre-programmed motor strategy and less of vestibular input while slow walking depends mainly on vestibular input so the sway in vestibular impairment is more in slowly walking than fast walking [32]. Patients with vestibular impairment make sway laterally, decrease walking speed and increase head movement as compensatory mechanisms [33]. Patients with unilateral vestibular impairment suffer from sway toward affected side [34].

The response of GVS is low when the patient is in standing but has a great effect during walking. This is an indicator that vestibular apparatus is very vital during walking more than during standing [35-40] GVS could stimulate the semicircular canal which evoked by angular acceleration and head velocity also could stimulate the static utricle and saccule which evoked by linear acceleration and head tilt [41].




Fig. 1. Underlying mechanisms of vestibular galvanic electric stimulation

The static and kinetic labyrinthine provide the CNS with successive sensory input about the linear and rotatory acceleration of the head which activates postural control during head movement via vestibulospinal and reticulospinal tracts [58].

GVS is an electric stimulation passed through application over mastoids producing modulation of the vestibular hair cells and their afferent activity [12,59]. GVS could decrease the abnormalities in walking performance especially in slow walking because the vestibular apparatus has a great role in postural stability in slow walking than in fast walking [60,61, 62,63,64].

Because the vestibular apparatus is a nonlinear fundamentally so the numbers of the neural network units increase the spread of stimulation producing large correct neural response due to more complicated neural network were involved in dynamic balance than in static balance which leads to improvement of stride length and time and walking velocity [65,66].

The vestibular disorder clinical picture includes static symptoms which include vestibular

nystagmus, head tilting, and body as a vestibulospinal sign, vertigo, perception sway and autonomic manifestations (nausea and vomiting). It needs short time for compensations to occur. Dynamic symptoms include impaired postural control and VOR deficits. It is poorly compensated and takes a long of time [67].

The vestibular restoration therapy is aiming for formation new learned dynamic strategies that can adjust posture during walking to reach the best performance of daily activities. The vestibule-plasticity depends on changing of the traditional neural circuits to well-developed strategies. The interplay between the brain plasticity and vestibule-plasticity in developing re-synaptic connection is the way for reaching the optimum dynamic balance control [68]. The molecular and cellular responses of the CNS due to feedback input and feedforward command as a result of physiotherapy training which produces increasing of neurotrophins, neurogenesis and new motor strategies which improve CNS plasticity [69].




Fig. 2. Underlying mechanisms of vestibular- plasticity mechanisms

The desensitization practices in vestibular restoration therapy is aiming for learning acquisition process to deal with difficult circumstances and responses could be used in vertigo treatment [74,76,77,78]. Vestibular restoration therapy should concentrate on adaptation mechanism for gaining the dynamic vestibular skill. Sensory substitution play a vital role in recovery of vestibular impairment by increasing the sensory feedback during close and open eyes, disturbance stimulus through different positions and manipulating the surface (unstable-rough-smooth-rubbery), upside down training. Proprioceptive training through static weight bearing and dynamic approximation plus a sense of weights to gain improvement of postural sway [73, 79.80,81] .

5. LIMITATION OF THE STUDY

- This study had specific limitations that can be concerned in the future research. Selection of larger groups of different forms and types of cerebral palsy children as (spastic diplegia and quadriplegia-dyskinesia-ataxia-mixed type of CP) are required to produce a foundation and principle when dealing with galvanic vestibular stimulation in pediatric rehabilitation.
- All of the hemiplegic CP children in this present study explore a sensory discomfort during application of galvanic vestibular stimulation.
- The GVS parameters in the present study were selected at the sensory threshold stimulus with the intensity of 0.5 mA intensity with long latency more than 200 ms for 30 minutes. Other current parameters and electrodes placement will be needed in further research to locate the effect of long run GVS on different cases in pediatric rehabilitation.

6. CONCLUSION

The combined vestibular training program and transmastoidal vestibular galvanic stimulation are suggested in improving walking performance and upright postural stability in a static and dynamic situation. So this selective physiotherapy approach may be used as a strong choice for improving walking and balance abilities in hemiplegic C.P children.

CONSENT

As per international standard written participant consent from the parents of the children studied has been collected and preserved by the authors.

ACKNOWLEDGEMENTS

I am grateful to my professor Emam El negmy for his help and advices.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Kerem Günel M, Livanelioğlu A. Cerebral palsy physiotherapy. Ankara, New Uzbek Press; 2009.
2. Berger W, Altenmueller E, Dietz V. Normal and impaired development of children's gait. *Hum Neurobiol.* 1984;3:163-170.
3. Liao HF, Jeng SF, Lai JS, Cheng CK, Hu MH. The relation between standing balance and walking function in children with spastic diplegic cerebral palsy. *Dev Med Child Neurol.* 1997;39:106-112.
4. Hubel DH, Wiesel TN. Binocular interaction in striate cortex of kittens reared with artificial squint. *J Neuro-Physiol.* 1965;28: 1041-59.
5. Merzenich MM. Soft wired: How the new science of brain plasticity can change your life. San Francisco, CA: Parnasus Publishing; 2013.
6. Merzenich MM, Van Vleet TM, Nahum. Brain plasticity-based therapeutics. *Front Hum Neurosci.* 2014;8:385. DOI: 10.3389/fnhum.2014.00385
7. Allred RP, Kim SO, Jonas. Use it and/or lose it—experience effects on brain remodeling across time after stroke. *Front Hum Neurosci.* 2014;8:379. DOI: 10.3389/fnhum.2014.00379
8. Hebb DO. The effects of early experience on problem-solving at maturity. *Am Psychol.* 1947;2:306-7.
9. Alwis DS, Rajan R. Environmental enrichment and the sensory brain: The role of enrichment in remediating brain injury. *Front Syst Neurosci.* 2014;8:156. DOI: 10.3389/fnsys.2014.00156
10. Cotman CW, Berchtold NC. Exercise: A behavioral intervention to enhance brain health and plasticity. *Trends Neurosci.* 2002;25:292-8.

- DOI: 10.1016/S0166-2236(02)02143-4
11. Lowenstein O. The effect of galvanic polarization on the impulse discharge from sense endings in the isolated labyrinth of the thornback ray (*Raja clavata*). *J Physiol.* 1955;127:104–117.
 12. Goldberg JM, Smith CE, Fernandez C. The relation between discharge regularity and responses to externally applied galvanic currents in the vestibular nerve afferents of the squirrel monkey. *J Neurophysiol.* 1984;51:1236–1256.
 13. Courjon JH, Precht W, Sirkin DW. Vestibular nerve and nuclei unit responses and eye movement responses to repetitive galvanic stimulation of the labyrinth in the rat. *Exp Brain Res.* 1987;66:41–48.
 14. Nardone A, Schieppati M. Balance in Parkinson's disease under static and dynamic conditions. *Mov Disord* 2006; 21(9):1515e20.
 15. Samoudi G, Nissbrandt H, Dutia MB, Bergquist F. Noisy galvanic vestibular stimulation promotes GABA release in the substantia nigra and improves locomotion in hemiparkinsonian rats. *PLoS One.* 2012;7(1):e29308.
 16. Blouin JS, Dakin CJ, van den Doel K, et al. Extracting phase-dependent human vestibular reflexes during locomotion using both time and frequency correlation approaches. *J Appl Physiol.* 2011;111: 1484e90.
 17. Dakin CJ, Inglis JT, Chua R, et al. Muscle-specific modulation of vestibular reflexes with increased locomotor velocity and cadence. *J Neurophysiol.* 2013;110: 86e94.
 18. Brandt T, Strupp M, Benson J. You are better off running than walking with acute vestibulopathy. *Lancet.* 1999;354:746.
 19. Schniepp R, Mohwald K, Wuehr M. Gait ataxia in humans: Vestibular and cerebellar control of dynamic stability. *J Neurol.* 2017;264:87e92.
 20. Schniepp R, Wuehr M, Neuhaeusser M, et al. Locomotion speed determines gait variability in cerebellar ataxia and vestibular failure. *Mov Disord.* 2012;27: 125e31.
 21. Matsuyama K, Drew T. Vestibulospinal and reticulospinal neuronal activity during locomotion in the intact cat. I. Walking on a level surface. *J Neurophysiol.* 2000;84: 2237e56.
 22. Mori S. Integration of posture and locomotion in acute de-cerebrate cats and in Awake, freely moving cats. *Prog Neurobiol.* 1987;28:161e95.
 23. Minor LB, Lasker DM, Backous DD, et al. Horizontal vestibule-ocular reflex evoked by high-acceleration rotations in the squirrel monkey. I. Normal responses. *J Neurophysiol.* 1999;82:1254e70.
 24. Sadeghi SG, Chacron MJ, Taylor MC, et al. Neural variability, detection thresholds, and information transmission in the vestibular system. *J Neurosci.* 2007;27: 771e81.
 25. Bent LR, Inglis JT, McFadyen BJ. When is vestibular information important during walking? *J Neurophysiol.* 2004;92: 1269e75.
 26. Ilg W, Giese MA, Gizewski ER, et al. The influence of focal cerebellar lesions on the control and adaptation of gait. *Brain.* 2008; 131:2913e27.
 27. Ilg W, Golla H, Thier P, et al. Specific influences of cerebellar dysfunctions on gait. *Brain.* 2007;130:786e98.
 28. Kavlak E, Ünal A, Tekin F, Altuğ F. Effectiveness of Bobath therapy on balance in cerebral palsy. *Cukurova Medical Journal.* 2018;43(4):975-981
 29. Kembhavi G, Darrah J, Magill-Evans J, et al. Using the berg balance scale to distinguish balance abilities in children with cerebral palsy. *Pediatr Phys Ther.* 2002; 14:92–99.
 30. Franjoine MR, Gunther JS, Taylor MJ. Pediatric balance scale: A modified version of the Berg Balance Scale for the school-age child with mild to moderate motor impairment. *Pediatr Phys Ther.* 2003;15: 114-28
 31. Goldberg JM, Fernandez C, Smith CE. Responses of vestibular nerve afferents in the squirrel monkey to externally applied galvanic currents. *Brain Research.* 1982;252:156-160. *Journal of Physiology.* 1999;517(3):931-939.
 32. Richard C. Fitzpatrick, Daniel L. Wardman, Janet L. Taylor effects of galvanic vestibular stimulation during human walking Prince of Wales Medical Research Institute, Sydney, Australia (Received 20 November 1998; accepted after revision 4; 1999.
 33. Glasauer S, Amorim MA, Vitte E, Berthoz A. Goal-directed linear motion in normal and labyrinthine-defective subjects.

- Experimental Brain Research. 1994;98: 323-335.
34. Halmagyi GM, Baloh RW. Overview of common syndromes of vestibular disease. In Disorders of the Vestibular System, ed. Baloh RW, Halmagyi GM. Oxford University Press, Oxford. 1996;291-299.
 35. Coates AC. Effect of varying stimulus parameters on the galvanic body-sway response. *Annals of Otology, Rhinology and Laryngology*. 1973;82:96-102.
 36. Nashner LM, Wolfson P. Influence of head position and proprioceptive cues on short latency postural reflexes evoked by galvanic stimulation of the human labyrinth. *Brain Research*. 1974;67:255-268.
 37. Smetanin BN, Popov KE, Gurfinkel VS, Shlykov VY. Effect of real and illusory movements on the human vestibule-motor reaction. *Neirofiziologiya*. 1988;20:250-255.
 38. Britton TC, Day BL, Brown P, Rothwell JC, Thompson PD, Marsden CD. Postural electro-myographic responses in the arm and leg following galvanic vestibular stimulation in man. *Experimental Brain Research*. 1993;94:143-151.
 39. Fitzpatrick R, Burke D, Gandevia SC. Task-dependent reflex responses and movement illusions evoked by galvanic vestibular stimulation in standing humans. *Journal of Physiology*. 1994;478:363-372.
 40. Inglis JT, Shupert CL, Hlavacka F, Horak FB. Effect of galvanic vestibular stimulation on human postural responses during support surface translations. *Journal of Neurophysiology*. 1995;73:896-90.
 41. Yang Y, Pu F, Lv X, Li S, Li J, Li D, et al. Comparison of postural responses to galvanic vestibular stimulation between pilots and the general populace. *Biomed Res Int*. 2015;567690.
 42. Day BL, Severac-Cauquil A, Bartolomei L, Pastor MA, Lyon IN. Human body-segment tilts induced by galvanic stimulation: A vestibular driven balance protection mechanism. *J Physiol*. 1997;500:661-672.
 43. Fitzpatrick R, McCloskey DI. Proprioceptive, visual and vestibular thresholds for the perception of sway during standing in humans. *J Physiol*. 1994;478:173-186.
 44. Lund S, Broberg C. Effects of different head positions on postural sway in man induced by a reproducible vestibular error signal. *Acta Physiol Scand*. 1983;117:307-309.
 45. Popov KE, Smetanin BN, Gurfinkel VS, Kudinova MP, Shlykov VYU. Spatial perception and vestibulomotor responses in man. *Neurophysiol*. 1986;18:548-553.
 46. Hlavacka F, Mergner T, Krizkova. Control of body vertical by vestibular and proprioceptive inputs. *Brain Res Bull*. 1996;40:431-435.
 47. Guerraz M, Thilo KV, Bronstein AM, Gresty MA. Influence of action and expectation on visual control of posture. *Cogn Brain Res*. 2001;11:259-266.
 48. Guerraz M, Day BL. Human body response to galvanic vestibular stimulation is not affected when the stimulus is selftriggered. *J Physiol*. 2001;531:142.
 49. Utz KS, Dimova V, Oppenlander K, Kerkhoff G. Electrified minds: Transcranial direct current stimulation (tDCS) and galvanic vestibular stimulation (GVS) as methods of non-invasive brain stimulation in neuropsychology review of current data and future implications. *Neuropsychologia*. 2010;48:2789e810.
 50. Stephan T, Deutschlander A, Nolte A, Schneider E, Wiesmann M, Brandt T, et al. Functional MRI of galvanic vestibular stimulation with alternating currents at different frequencies. *Neuroimage*. 2005; 26:721e32.
 51. Moliadze V, Atalay D, Antal A, Paulus W. Close to threshold transcranial electrical stimulation preferentially activates inhibitory networks before switching to excitation with higher intensities. *Brain Stimul*. 2012;5:505e11.
 52. Uchino Y, Kushiro K. Differences between otolith- and semicircular canal activated neural circuitry in the vestibular system. *Neurosci Res*. 2011;71:315e27.
 53. Lowrey CR, Bent LR. Modulation of the soleus H-reflex following galvanic vestibular stimulation and cutaneous stimulation in prone human subjects. *Muscle Nerve*. 2009;40:213e20.
 54. Kennedy PM, Inglis JT. Modulation of the soleus H-reflex in prone human subjects using galvanic vestibular stimulation. *Clin Neurophysiol*. 2001;112: 2159e63.
 55. Roerdink M, Hlavackova P, Vuillerme N. Effects of plantar-flexor muscle fatigue on the magnitude and regularity of center-of-pressure fluctuations. *Exp Brain Res*. 2011;212:471e6.
 56. Hlavackova P, Pradon D, Vuillerme N. Control of bipedal posture following localized muscle fatigue of the plantar-

- flexors and finger-flexors. *Eur J Appl Physiol.* 2012;112:789e93
57. Cathers I, Day BL, Fitzpatrick RC. Otolith and canal reflexes in human standing. *J Physiol.* 2005;563:229e34.
 58. Baloh RW, Honrubia V. *Clinical neurophysiology of the vestibular system.* New York: Oxford University Press; 2001.
 59. Aw ST, Todd MJ, Aw GE, et al. Gentamicin: Vestibule-toxicity impairs human electrically evoked vestibular-ocular reflex. *Neurology.* 2008;71:1776e82.
 60. Fujimoto C, Yamamoto Y, Kamogashira T, et al. Noisy galvanic vestibular stimulation induces a sustained improvement in body balance in elderly adults. *Sci Rep.* 2016;6:37575.
 61. Iwasaki S, Yamamoto Y, Togo F, et al. Noisy vestibular stimulation improves body balance in bilateral vestibulopathy. *Neurology.* 2014;82:969e75.
 62. Iwasaki S, Karino S, Kamogashira T, et al. Effect of noisy galvanic vestibular stimulation on ocular vestibular-evoked myogenic potentials to bone conducted vibration. *Front Neurol.* 2017;8:26.
 63. Wuehr M, Nusser E, Decker J, et al. Noisy vestibular stimulation improves dynamic walking stability in bilateral vestibulopathy. *Neurology.* 2016;86:2196e202.
 64. Wuehr M, Nusser E, Krafczyk S, et al. Noise-enhanced vestibular input improves dynamic walking stability in healthy subjects. *Brain stimulation.* 2016;9:109e16.
 65. Collins JJ, Chow CC, Imhoff TT. Stochastic resonance without tuning. *Nature.* 1995;376:236e8.
 66. Collins JJ, Imhoff TT, Grigg P. Noise-enhanced information transmission in rat SA1 cutaneous mechanoreceptors via aperiodic stochastic resonance. *J Neurophysiol.* 1996;76:642e5.
 67. Halmagyi GM, Curthoys IS, Cremer PD, Henderson CJ, Todd MJ, Staples MJ, et al. The human horizontal vestibulo-ocular reflex in response to high-acceleration stimulation before and after unilateral vestibular neurectomy. *Exp Brain Res.* 1990;81:479.
DOI: 10.1007/BF02423496
 68. Lacour M, Demanze LB. Interaction between vestibular compensation mechanisms and vestibular rehabilitation therapy: 10 recommendations for optimal functional recovery *Frontiers in Neurology* published; 2015.
DOI: 10.3389/fneur.2014.00285
 69. Cotman CW, Berchtold NC. Exercise: Behavioral intervention to enhance brain health and plasticity. *Trends Neurosci.* 2002;25:292–8.
DOI: 10.1016/S0166-2236(02)02143-4
 70. Travo C, Gaboyard-Niay S, Chabbert C. Plasticity of Scarpa's ganglion neurons as a possible basis for functional restoration within vestibular end organs. *Front Neurol.* 2012;3:91.
DOI: 10.3389/fneur.2012.00091
 71. Vande Berg R, Guinand N, Stokroos R, Guyo JP, Kingma H. The vestibular implant: Quovadis? *Front Neurol.* 2011;2:47.
DOI: 10.3389/fneur.2011.00047
 72. Borel L, Lopez C, Peruch P, Lacou RM. Vestibular syndrome: A change in internal spatial representation. *Clin Neurophysiol.* 2008;38:375–89.
DOI: 10.1016/j.neucli.2008.09.002
 73. Horak FB. Postural compensation for vestibular loss. *Resto Neurol Neurosci.* 2010;28:57–68.
DOI: 10.3233/RNN-2010-0515
 74. Norré ME, Becker AM. Vestibular rehabilitation training. Specificity of adequate exercises. *Arch Otolaryngol Head Neck Surg.* 1988;114:883–6.
DOI: 10.1001/archotol.1988.0186020006702
 75. Brandt T, Daroff RB. Physical therapy for benign paroxysmal positional vertigo. *Arch Otolaryngol.* 1980;106:484–5.
DOI: 10.1001/archotol.198000790320036009
 76. Brandt T, Daroff RB. Physical therapy for benign paroxysmal positional vertigo. *Arch Otolaryngol.* 1980;106:484–5.
DOI: 10.1001/archotol.19800079032003600976
 77. Tee LH, Chee NW. Vestibular rehabilitation therapy for the dizzy patient. *Ann Acad Med Singapore.* 2005;34:289–94.
 78. Bronstein AM, Golding FJ, Gresty MA. Vertigo and dizziness from environmental motion: Visual vertigo, motion sickness, and drivers' disorientation. *Semin Neurol.* 2013;33:219–30.
DOI: 10.1055/s-0033-1354602
 79. Bittar RS, Barros CdeG. Vestibular rehabilitation with biofeedback in patients with central imbalance. *Braz J Otorhinolaryngol.* 2011;77:356–61.
DOI: 10.1590/S1808-86942011000300014
 80. Dozza M, Chiari L, Hlavacka F, Cappello A, Horak FB. Effects of linear versus sigmoid coding of visual or auditory

biofeedback for the control of upright stance. *EEE Trans Neural Syst Rehabil Eng.* 2006;14:505–12.
DOI: 10.1109/TNSRE.2006.886732

81. Dozza M, Horak FB, Chiari L. Auditory biofeedback substitutes for loss of sensory information in maintaining stance. *Exp Brain Res.* 2007;178:37–48.
DOI: 10.1007/s00221-006-0709-y

© 2019 Azzam; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48264>